

## **BIMATOPROST E/D**

**CLASS:** Ophthalmic Agent, Antiglaucoma; Prostaglandin, Ophthalmic

**INDICATIONS:** Reduction of intraocular pressure (IOP) in patients with open-angle glaucoma or ocular hypertension; hypotrichosis treatment of the eyelashes

### **AVAILABLE DOSAGE FROM THE HOSPITAL:**

BIMATOPROST 0.3MG/ML EYE DROP

### **TRADE NAMES:**

### **DOSAGE:**

- **Dosing: Adult**

Open-angle glaucoma or ocular hypertension: Ophthalmic: Instill 1 drop into affected eye(s) once daily in the evening; do not exceed once-daily dosing (may decrease IOP-lowering effect). If used with other topical ophthalmic agents, separate administration by at least 5 minutes.

Hypotrichosis of the eyelashes: Ophthalmic, topical: Place one drop on applicator and apply evenly along the skin of the upper eyelid at base of eyelashes once daily at bedtime; repeat procedure for second eye (use a clean applicator).

- **Dosing: Geriatric**

Refer to adult dosing

### **COMMON SIDE EFFECT:**

**>10%:** Ocular: Conjunctival hyperemia (25% to 45%; Latisse®: <4%), growth of eyelashes, ocular pruritus (>10%; Latisse®: <4%)

### **PREGNANCY RISK FACTORS:** C