

DORZOLAMIDE EYE DROP

CLASS: Carbonic Anhydrase Inhibitor (Ophthalmic); Ophthalmic Agent, Antiglaucoma

INDICATIONS: Treatment of elevated intraocular pressure in patients with ocular hypertension or open-angle glaucoma

AVAILABLE DOSAGE FROM THE HOSPITAL:

DORZOLAMIDE 2% EYE DROP, DORZOLAMIDE 2% EYE DROP, DORZOLAMIDE HCL 2%+ TIMOLOL 0.5% E/D

TRADE NAMES:

DOSAGE:

- **Dosing: Adult**

Reduction of intraocular pressure: Ophthalmic: Instill 1 drop in the affected eye(s) 3 times/day

- **Dosing: Geriatric**

Refer to adult dosing.

- **Dosing: Renal Impairment**

Clcr < 30 mL/minute: Use is not recommended (has not been studied).

- **Dosing: Hepatic Impairment**

No dosage adjustment provided in manufacturer's labeling (has not been studied); use with caution.

COMMON SIDE EFFECT:

1% to 10%:

Gastrointestinal: Bitter taste following administration (25%)

Ocular: Burning, stinging or discomfort immediately following administration (33%); superficial punctate keratitis (10% to 15%); signs and symptoms of ocular allergic reaction (10%)

PREGNANCY RISK FACTORS: C