

## **OXYBUTININE**

**Class:** Antispasmodic Agent, Urinary

**Indications:** Treatment of symptoms associated with overactive uninhibited neurogenic or reflex neurogenic bladder (eg, urgency, frequency, leakage, urge incontinence, dysuria); treatment of symptoms associated with detrusor overactivity due to a neurological condition (eg, spina bifida) (extended release tablet only)

**Available dosage form in the hospital:**

OXYBUTININE 5MG TAB

**Dosage:**

**-Overactive bladder:**

*Oral:*

-Immediate release: 5 mg 2-3 times daily; maximum: 5 mg 4 times daily

-Extended release: Initial: 5-10 mg once daily, adjust dose in 5 mg increments at weekly intervals; maximum: 30 mg once daily

**Geriatric**

*Oral:* Immediate release: Initial: 2.5 mg 2-3 times daily; increase cautiously

**Renal Impairment:**

No dosage adjustment provided in the manufacturer's labeling (not studied); use with caution.

**Hepatic Impairment:**

No dosage adjustment provided in the manufacturer's labeling (not studied); use with caution.

**Common side effect:**

**Oral:**

Central nervous system: Dizziness, drowsiness

Gastrointestinal: Xerostomia ( dose related), constipation, nausea

**Pregnancy Risk Factor:** B