

## **PREDNISOLONE ACETATE 0.12% ,1% EYE DROPS**

**CLASS:** Corticosteroid, Ophthalmic

**INDICATIONS:** Treatment of palpebral and bulbar conjunctivitis; corneal injury from chemical, radiation, thermal burns, or foreign body penetration; steroid-responsive inflammatory ophthalmic diseases

### **AVAILABLE DOSAGE FROM THE HOSPITAL:**

PREDNISOLONE 20MG TAB, PREDNISOLONE 50MG TAB, PREDNISOLONE 5MG TAB, PREDNISOLONE ACETATE 0.12% EYE DROPS, PREDNISOLONE ACETATE 1% EYE DROPS, PREDNISOLONE +NEOMYCIN +POLYMYXINE -B EYE /DROPS

### **DOSAGE:**

**Conjunctivitis:** Ophthalmic (suspension/solution): Instill 1-2 drops in the eye 2-4 times daily

### **Geriatric**

Refer to adult dosing.

**COMMON SIDE EFFECT:** Frequency not defined.

Ocular: Conjunctival hyperemia, conjunctivitis, corneal ulcers, delayed wound healing, glaucoma, intraocular pressure increased, keratitis, loss of accommodation, optic nerve damage, mydriasis, posterior subcapsular cataract formation, ptosis, secondary ocular infection

**PREGNANCY RISK FACTORS:** C