

SOD.HYALURONATE FOR INTERSTITIAL CYSTITIS

Class: Antirheumatic, Miscellaneous; Ophthalmic Agent, Viscoelastic; Skin and Mucous Membrane Agent, Miscellaneous

Indications: Use - Unlabeled

Treatment of refractory interstitial cystitis

Available dosage form in the hospital: SOD.HYALURONATE 0.8MG/ML FOR INTERSTITIAL CYSTITIS (50ML VIAL)

Dosage:

Interstitial cystitis, refractory (unlabeled use): Intravesical (unlabeled route): 40 mg in 50 mL saline intravesically (retain in bladder for at least 30 minutes) once weekly for 4 weeks, then monthly for up to 1 year in patients showing an initial response

Common side effect:

Frequencies and/or type of local reaction may vary by formulation and site of application/injection.

-Local: Injection site (intra-dermal): Erythema (75% to 93%), tenderness (61% to 92%), swelling (81% to 91%), pain (47% to 90%), firmness (86% to 89%), bruising (52% to 87%), lumps/bumps (56% to 83%), skin discoloration (33% to 78%), pruritus (25% to 36%)

-Neuromuscular & skeletal: Arthralgia (intra-articular 25%)

Pregnancy Risk Factor: Adverse events were not observed in animal reproduction studies. Safety for use in pregnant women has not been established.