

**•SODIUM CROMOGLYCATE NEBULIZER 20MG**

**Class:** Mast Cell Stabilizer

**Indications:**

Adjunct in the prophylaxis of allergic disorders, including asthma; prevention of exercise-induced bronchospasm.

**Available dosage form in the hospital:**

-SODIUM CROMOGLYCATE NEBULIZER 20MG

**Dosage:**

**-Asthma:***Nebulization solution:* Initial: 20 mg 4 times/day; usual dose: 20 mg 3-4 times/day

**-Prophylaxis of bronchospasm (allergen- or exercise-induced):**

*Nebulization solution:* Single dose of 20 mg.

**Geriatric**

Refer to adult dosing.

**Renal Impairment:**

Specific guidelines not available; consider lower dose of oral product.

**Hepatic Impairment:**

Specific guidelines not available; consider lower dose of oral product.

**Common side effect:**

Frequency not defined.

- Cardiovascular: Angioedema, chest pain, edema, flushing, palpitation, premature ventricular contractions, tachycardia
- Central nervous system: Anxiety, behavior changes, convulsions, depression, dizziness, fatigue, hallucinations, headache, irritability, insomnia, lethargy, migraine, nervousness, hypoesthesia, postprandial lightheadedness, psychosis
- Dermatologic: Erythema, photosensitivity, pruritus, purpura, rash, urticaria
- Gastrointestinal: Abdominal pain, constipation, diarrhea, dyspepsia, dysphagia, esophagospasm, flatulence, glossitis, nausea, stomatitis, vomiting
- Genitourinary: Dysuria, urinary frequency
- Hematologic: Neutropenia, pancytopenia, polycythemia

- Hepatic: Liver function test abnormal
- Local: Burning
- Neuromuscular & skeletal: Arthralgia, leg stiffness, leg weakness, myalgia, paresthesia
- Otic: Tinnitus
- Respiratory: Dyspnea, pharyngitis
- Miscellaneous: Lupus erythematosus

**Pregnancy Risk Factor: B**